

Adam Zack

From: Lynda Guernsey
Sent: Thursday, July 8, 2021 9:03 AM
To: Comp Plan Update
Subject: FW: Ferry Limitations & Land Use Redesignations = Health Impacts
Attachments: Wetland_Function&Loss.pdf; Wetland_Function&Loss.pdf

Please see email below and attachments.

Lynda Guernsey, Administrative Specialist II – Direct Line (360) 370-7579
SAN JUAN COUNTY
DEPARTMENT OF COMMUNITY DEVELOPMENT
(360) 378-2354 | 135 Rhone Street | PO Box 947 | Friday Harbor, WA 98250

From: Jan Alderton <janetmalderton@gmail.com>
Sent: Wednesday, July 7, 2021 6:48 PM
To: Ingrid Gabriel <ingridg@sanjuanco.com>; Cindy Wolf <cindyw@sanjuanco.com>; Jamie Stephens <jamies@sanjuanco.com>; Christine Minney <christinem@sanjuanco.com>; Lynda Guernsey <LyndaG@sanjuanco.com>
Subject: Ferry Limitations & Land Use Redesignations = Health Impacts

CAUTION: This email originated from outside of the organization. Do not click links or open attachments unless you recognize the sender and know the content is safe.

Ferry Limitations & Land Use Redesignations = Health Impacts

Dear Planning Commissioners and County Councilors,

For the San Juan Islands, declining ferry services are more than an inconvenience. Many older residents, including myself, have increasing healthcare needs that cannot be addressed locally. For this large segment of our community, ferry service to the mainland is truly an essential public service.

The Growth Management Act requires that essential public services, including transportation infrastructure, be adequate to accommodate projected growth. Redesignating a large acreage of Forest Resource Lands to Rural Farm Forest could therefore violate the Growth Management Act.

Until ferry service is significantly improved, please do not increase the build-out potential in our islands. Retired ferry captain Ken Burtness, who serves on the Ferry Advisory Committee, told me recently that this will take a minimum of ten years.

The Seattle Times article, below, describes the perfect storm of factors that have impacted our ferry service. These problems will only get worse over time unless significant new sources of funding are found.

Ferry funding took a huge hit after passage of the Tim Eyman-sponsored Initiative 695 that limited car tab fees in 2000. The resulting reduction in funding severely impacted our Washington State Ferry system. Fare increases alone are not "the solution". Even with increased funding, the timeline for improvement is a minimum of ten years. The scope of the work includes the replacement of aging ferries with new vessels and adequate provision of reserve ferries (that are called into service when a ferry requires major repairs).

The problems associated with insufficient crew numbers for the ferries will only be addressed by significantly increased funding of the ferries. Increased salaries and changes to on-call working requirements will be necessary to attract people to a career dedicated to working on our ferries.

How Are Ferry Capacity and Redesignation of Forest Resource Lands Connected?

Redesignation of Forest Resource Lands to Rural Farm Forest will ultimately add to the "build out" potential in our islands and therefore to the number of people using the ferries. Even though the "development potential" for the redesignated parcels is theoretically not increased in this proposal, history tells me that the allowed density per parcel will change. Guest houses would be allowed and the percentage of allowable impervious surfaces would increase. Individual parcel owners will inevitably appeal to increase the density to one residence per five acres plus the potential guest house. And appeals will be granted because the prevailing attitude in our islands is that "one more" development project will have minimal impacts. The problem is that cumulative "one more" impacts become truly significant. This is like Continental Drift. The rate of change is small each year, but over time, the magnitude of change is dramatic.

There are also environmental reasons to oppose redesignations that will increase the build-out potential in our islands. Reductions in our forest

lands will take us in the opposite direction needed to address Climate Change. Trees remove carbon dioxide from the atmosphere and trees also help to recharge our groundwater supplies. Removal of trees and replacement of forests with the impervious surfaces that are part of residential development leads to increased stormwater flows that are not trivial to mitigate. Mitigation of impacts never fully replaces the ecosystem services of the natural habitat that is destroyed. I can supply scientific articles that clearly demonstrate this "Myth of Mitigation". See the attached journal article about how wetlands "restoration" does not fully replace the lost functions and values of natural wetlands,

Please do not approve the Docket Proposal that would redesignate Forest Resource Lands to Rural Farm Forest.

Janet Alderton

Orcas Island

https://www.seattletimes.com/seattle-news/transportation/why-your-ferry-might-be-late-or-canceled-this-summer/?utm_source=marketingcloud&utm_medium=email&utm_campaign=Morning+Brief+7-7-2021_7_7_2021&utm_term=Active%20subscriber

Why your ferry might be late — or canceled — this summer

July 7, 2021 at 6:00 am

A shortage of boats and crew at Washington State Ferries is subjecting travelers to a summer of cancellations.

A high number of trips have already been canceled this year, according to WSF spokesperson Ian Sterling. Since February, at least 57 round-trip sailings have been called off due to staffing shortages. There were only 10 and five cancellations in the same time period respectively in 2019 and 2018.

As ridership is returning to pre-pandemic levels, passengers are starting to notice lapses in service, Sterling said. For an agency that typically completes [99% of its scheduled trips](#), those cancellations and travel alerts have been demoralizing for staff members, he said. Some trips were canceled in the San Juan Islands over the weekend, and the Bremerton-Seattle route Tuesday morning.

For every cancellation, Sterling said there are 10 more that were almost canceled. Dispatch scrambled and employees, scheduled to be off, volunteered to come in.

It's resulting in burnout among staff, said Jay Ubelhart, president of the Inlandboatmen's Union.

A crew shortage knocked the Port Townsend-Coupeville route to just one ferry at the height of touring season. Chief of Staff Nicole McIntosh blamed it on "an unprecedented staffing challenge" [in a personnel letter posted on all vessels](#) June 24.

While thanking crew for extraordinary work to keep boats moving in the pandemic, she urged them to answer last-minute calls to work. "We have an obligation to the taxpayers of the state to not miss sailings due to crewing. When we do it lets down thousands of customers who trust and depend on us to get them safely to their destinations," she wrote.

The agency reports nearly 300,000 riders during the holiday weekend. Saturday was the highest ridership day with nearly 86,000 riders, the most in a single day since Sept. 21, 2019, according to Sterling.

Though fleet size has improved since May, when WSF was short three boats, the usual summer fleet of 19 vessels is down to 18 after an engine fire this spring aboard the ferry Wenatchee will [keep that boat docked for months](#).

That's barely manageable, because pandemic border restrictions have canceled service from Anacortes to Sidney, B.C., freeing up one boat for domestic trips.

Depending on the vessel and the number of passengers, the U.S. Coast Guard mandates a certain number of crew members before a boat can leave the dock.

Before 2012, crews operated with an extra member in case someone calls in sick, or has car trouble on the way to work, according to Sterling. To save money, ferries are now scheduled with the minimum number of crew members. One absent colleague can result in a canceled sailing — forcing passengers to wait for the next trip.

Washington State Ferries recommends travelers use its [travel and traffic guide](#), planning trips [early in the morning](#) or later at night and checking the [travel alerts bulletin](#) for schedule changes. If there is a long wait, WSF also suggests driving to another route without a delay.

Difficulties recruiting

The maritime industry in general struggles with recruiting young adults. But that has especially slowed during the pandemic for WSF, which faces a coming wave of retirements.

Working on a ferry is "a total shock to the system" compared to a typical 9-to-5 office job, said Ubelhart, who has worked on the state's ferries for 20 years. To even apply for the most junior position, people must complete at least 14 days of training and often pay up to \$400 out of pocket, he said.

Work schedules are unpredictable both on a day-to-day and season-to-season basis, especially for new employees without seniority or a steady shift, he said. New employees usually start out on call, and wait for a dispatcher to call with work.

“They’ll call you any time and you’re expected to jump into your uniform, grab a lunch and go to work,” Ubelhart said.

While working for WSF offers an upward career trajectory and a pension, Sterling said the agency has a hard time attracting young adults when competing with Seattle tech companies and the changing nature of work.

During the pandemic, Kitsap County resident Elliott Smith quit his office job to complete maritime training, with an eye toward work on a ferry. Instead, he began this summer working on a cargo ship in Beaumont, Texas.

Among other reasons, Smith was unwilling to spend at least two years on call, where he might be told to drive to any ferry terminal on any given day, or maybe not work at all.

“I would be applying for Washington State Ferries today, if I knew I could get a steady paycheck, but I’m not going to gamble for two years,” Smith said.

Wages also have not kept up with Seattle’s rising cost of living, said Ryan Brazeau, a fourth-generation ferry worker and officer in the Inlandboatmen’s Union. Tech has taken over the downtown area and people are unwilling to make long commutes to work for WSF, he said.

“This career is not built for a family either,” he said. “During this year COVID is taking a toll on people that have kids.”

As a taxpayer and Puget Sound-area native, Smith said it bothers him that the ferry logistics seem so disorganized. “We buy these multimillion-dollar boats, we keep them in shape, then we can’t run them because we don’t have enough people,” Smith said by phone from Texas last month.

Coming retirements and open positions

Since July 2019, 29 mates and captains have retired and 75 seamen have left, according to WSF. While the agency says it has managed to recruit enough seamen, they are still short about 10 senior deck crew.

The most recent breaking point has been the engine room, where around 34 employees have left, according to WSF. But a new class of about seven oilers is about to graduate, Sterling said, taking some pressure off the engine room staffs.

The agency is also anticipating more retirements. At least half the remaining senior engineers and deck crew are eligible for retirement, he said.

WSF couldn’t afford to build new boats for a decade, due to fallout from a car-tab cut in 2000. New taxes since the mid-2010s are replenishing the budget. The next boat, a 144-car hybrid

electric-diesel vessel, will [begin construction soon at Vigor in Seattle](#), but won't sail until 2024. The existing fleet is working past retirement age, and three vessels are at least 60 years old.

"Any time a ferry goes down, it shuts the whole system," Brazeau said.

Ferry careers

Ordinary seaman: Responsible for cleaning, first aid, firefighting, lookout. Requires 13 days training. Average pay, \$22.56/hour.

Able-bodied seaman: Responsible for vehicle loading, lifeboats, knot tying, assisting mates/captains. Requires two years experience, nine days of school. \$31.85/hour.

Mate: Shares navigation with captain, supervises vehicle loading, crowd management. Requires four years sea time, 33 days of officer school, 163 days of pilotage, study, training. \$44.62/hour.

Captain: Full command of vessel and passenger safety. Requires five to six years sea time, 35 days study, Coast Guard master's license. \$55.27/hour.

Oilers: Responsible for inspection of all systems, operating equipment, repair fixtures, mechanical and electrical aide for repairs. Requires 14 days of training. Average pay \$28.49/hour.

Assistant engineers: Responsible for running, operation and maintenance of the propulsion and electrical systems aboard the vessels. Average pay \$42.74/hour.

Chief engineers: Responsible for implementing all federal and state regulations, WSF policies and procedures, and any directives from WSF management. Average pay \$50.73/ hour.

Staff chief engineer: Responsible for all standard maintenance on the vessel, reviews and approves all maintenance recommendations for improvements on the vessel. Average pay \$66.77/hour.




Figure 2. Recovery trajectories of animal and plant richness and density. Chronosequences of the means (\pm standard error [SE]) of the response ratios (see Materials and Methods) of restored or created wetlands at successive age classes of 5 y or 10 consecutive y for vertebrates and macroinvertebrates density and richness (A) and for plant density and richness (B). Insufficient data points meeting our plotting criteria (see Materials and Methods) were available to plot for macroinvertebrate richness. The zero value dashed line represents reference wetlands (N, number of data points used to calculate the mean [\pm SE] per age class; Y, years after restoration). Subscripts are as follows: md, macroinvertebrates density; pd, plant density; pr, plant richness; vd, vertebrate density; vr, vertebrates richness). doi:10.1371/journal.pbio.1001247.g002

restoration were not plotted in Figure 1A because the sample size did not meet our criteria for average points, see Materials and Methods section, on this graph). Since phosphorus storage appeared only slightly changed, the overall lack of recovery of biogeochemical functioning may have been driven largely by the low recovery of the carbon storage and the low accumulation of soil organic matter (see Text S1).

Effects of Size and Environmental Setting

Comparing wetland recovery trajectories under different conditions may shed light on factors that impede or facilitate recovery. Although biogeochemical responses in both restored and created wetlands were similar, biological structure in created wetlands approached reference conditions more quickly (Figure S3A and S3B; Table S5). Created wetlands may have been engineered to force the initial system towards defined reference conditions [39].

Ecosystem size and local and regional context affect wetland recovery. Large wetlands (>100 ha) appeared to recover their biological structure and biogeochemical functions sooner after restoration or creation than smaller wetlands (Figures 3 and S4; Table S4; data from 13 studies of 25 wetlands). This differential recovery suggests that small wetlands may not provide adequate local resources or connectivity for local biota to restore preimpact functioning. Restored and created wetlands, particularly if small, may have become more isolated and surrounded by more fragmented landscapes than they had been before impact [40]. Also, small wetlands would only be able to support a limited number of individuals, and thus, will not be able to support all the species, particularly taxa with large body sizes, formerly capable of occupying the area [41].

Regional climate had a strong effect on the sequence and rate of wetland recovery following restoration. As expected, warm temperatures accelerate ecosystem processes [7,13,42], including those mediating biological and biogeochemical recovery after wetland restoration or creation. In tropical and summer-warm temperate climates, wetlands approached reference conditions relatively rapidly, while wetlands restored in cold climates had not recovered to reference conditions after 50 y (Figure 4A and 4B; Tables S3 and S5). In tropical climates only, biogeochemical

variables recovered to reference levels before biological structure did (data from eight studies of eight wetlands). Whether this difference in recovery sequence is a real aspect of tropical wetlands, or an artifact of small sample size, remains to be seen. In a much larger sample of studies from temperate climates this sequence was reversed, and biogeochemical recovery was slower. Biological structural variables appeared recovered 5 y after restoration, while even 30 y after restoration, biogeochemical functions had only recovered to 79% of reference levels (data from 83 studies of 302 wetlands). In cold climates, corresponding biogeochemical recovery was only 53% 50 y after restoration; both biogeochemical functions and biological structure variables

Figure 3. Effect of size on wetland recovery. Evolution of the mean (\pm standard error [SE]) of the response ratios (see Materials and Methods) of restored or created wetlands at successive size categories for wetlands between 0 y to 5 y after restoration or creation. The zero value dashed line represents reference wetlands. Mean (\pm SE) at 0.1 ha was estimated for wetlands with sizes \leq 0.1 ha. Means (\pm SE) at 1 ha were estimated for wetlands in which sizes ranged between 0.1 ha and 1 ha. The same approach was used to estimate the means (\pm SE) at 10, 100, 1,000, and 10,000 ha (N, number of data points used to calculate the mean [\pm SE] per age class; size, size in hectares of the restored wetlands. Subscripts are as follows: bp, biogeochemical processes; bs, biological structure). doi:10.1371/journal.pbio.1001247.g003




Figure 4. Effects of climate and hydrology on wetland recovery trajectories. Chronosequences of the means (\pm standard error [SE]) of the response ratios (see Materials and Methods) of restored and created wetlands at successive age classes of 5 y or 10 consecutive y for biogeochemical functions and for biological structures under contrasting climates (A and B), and under different hydrologic connectivity (C and D) [31]. The zero value dashed line represents reference wetlands. The arrow (B) indicates the outlier mean value of two restoration studies with extremely low recovery rates (N, number of data points used to calculate the mean [\pm SE] per age class; Y, years after restoration). Subscripts are as follows: bp, boreal peatland; d, depressionnal; hc, humid cold; ht, humid temperate; r, riverine; str, seasonal tropical; ste, seasonal temperate; t, tidal. doi:10.1371/journal.pbio.1001247.g004

remained statistically distinct from reference conditions for the entire (50-y) chronosequence (Figure 4A and 4B; Tables S3 and S5; data from 33 studies of 311 wetlands).

Hydrologic setting [43] also affected recovery (Figure 4C and 4D; Tables S3 and S5). Riverine and tidal wetlands, linked to larger hydrologic regimes by natural flow variation, recovered biogeochemical functions and biological structure after 20 y and 30 y, respectively (data from 73 studies of 210 wetlands). These results are similar to those (15 y to 25 y to recover the original biotic composition and diversity) found by Borja et al. [8] in 51 globally distributed estuarine and coastal ecosystems. In contrast, wetlands in inland depressions that were watered by precipitation or groundwater flow had not recovered to reference conditions even after 50 y following restoration (data from 36 studies of 358 wetlands). Peatlands (usually only the upper layer [<1 m] of peat was removed) recovered biological structure immediately, but 30 y after restoration, biogeochemical functioning in peatlands remained statistically lower than in reference wetlands (data from 11 studies of 18 wetlands).

Slow Recovery or Alternative States?

Two hypotheses could explain the lag in biological and biogeochemical recovery of the biological structure and biogeo-

chemical functioning. First, the chronosequences we examined may be too short (<30 y) for full recovery, especially of carbon and nitrogen storage [44]. Second, restored wetlands may have shifted to alternative states, different from their condition before degradation [14,15]. The subreference plateaus of soil organic accumulation, carbon storage, and general biogeochemical functioning could support the second hypothesis of alternative states in restored systems. Slow recovery of plant density and richness might be linked to lags in carbon storage. Mutualist symbionts critical for plant productivity (e.g., N-fixing bacteria [2] or mycorrhizal fungi [45]) may be absent in recently (<50 y) restored wetland soils. Alternatively, fast-growing, early successional terrestrial plants, and potentially also wetland plants, usually allocate most of their carbon to photosynthetically active structures of low density and high nutrient content, which are easily grazed or rapidly decomposed, retarding local storage of carbon [46,47].

Comparison with Other Findings

Two other studies have assessed recovery rates of large scale natural ecosystems following disturbance or perturbations [7,12]. Both of these studies examined a broad range of ecosystem types (terrestrial, freshwater, and marine), including wetlands. Jones and

Schmitz [12] found that across ecosystems and perturbation types (natural and human-caused), about half of the tracked response variables were considered by original authors to have recovered to preimpact states. Jones and Schmitz computed averaged recovery times for the subsamples of variables and cases that primary authors considered to have recovered over the course of their studies. These recovery times ranged from about 10 y to 40 y, and were longer for forests, and following human-caused, rather than natural perturbations. To assess whether systems had recovered or not, Jones and Schmitz used authors' expert opinion, return to historic initial conditions, or approach to parallel reference states (our study evaluated recovery only for studies using the last of these criteria). Given the narrower scope of our study (assessing wetlands only), and our different analysis approach, estimated recovery times from these two reviews are surprisingly similar. Rey Benayas et al. [7] studied recovery across a wide range of human-perturbed ecosystems, including wetlands. Using (as we did) the response ratio of restored to reference ecosystems, Benayas et al. found biodiversity and selected ecosystem services to be 86% and 80% recovered in a sample of 89 cases pooled over all age categories since perturbation. Interestingly, they reported slightly (6%) higher recovery in biological variables compared to ecosystem services (nutrient cycling; primary production; provisioning of timber, fish, and food crops; and regulation of climate, water supply, and soil). These ecosystem services overlap in part with categories of biogeochemical variables in our study (e.g., carbon and nutrient storage and cycling). The similarity between their results and our finding (that biological variables were 9% more recovered than these biogeochemical responses) suggests that structural recovery might often be necessary to achieve functional recovery.

Conclusions

Our meta-analysis suggests that recovery of wetlands following restoration as currently practiced is often slow and incomplete. In warm climates, and in settings linked to riverine or tidal flows, recovery may proceed more rapidly. Recovery may also be more likely and more rapid if >100 contiguous ha are restored. In many wetlands, however, ecosystem services may not be fully recovered even when wetlands appear to be biologically restored. If markets for ecosystem services and mitigation offsets from restored or created wetlands are used to justify further wetland degradation, net loss of global wetland services will continue and likely accelerate (see also Race and Fonseca [48]). We join other wetland ecologists and restoration scientists in calling for better scientific understanding of biotic and abiotic factors that constrain ecosystem restoration. For our common future, we need more realistic, long-term evaluations to find better ways to alleviate constraints limiting the recovery of wetland ecosystems.

Materials and Methods

Literature Search

On the 22nd of December 2010 a reference search was done in the scientific database ISI Web of Science – SCI-Expanded. The terms used were “(wetland* or floodplain* or peatland* or marsh* or mangrove*) same (restor* or creat* or re-creat* or rehabilit*)”. We used these terms to cover a wide variety of wetlands as defined in the Article 1.1 of the Ramsar Convention text [7]. For this analysis, we considered restored wetlands to be wetlands recreated on sites where wetlands had formerly existed but been drained or otherwise severely degraded. Created wetlands were described by authors as wetlands built on sites that lacked previous wetland history. We selected studies of wetlands under natural hydrological regimes, planted with native species, and in which no allochtho-

nous substrates were imported during the restoration or creation activities. For this reason, the term “construct*” was not included in the search terms, because we found in an independent search that >99% of the studies of constructed wetlands were of highly artificial systems not maintained under natural conditions. The search produced 2,959 selected articles. We applied the general selection criterion: “Articles must compare measurements of structural components and biogeochemical processes in restored or created and reference wetlands at a known age.” Under this criterion we selected 172 articles. These articles were read, and those in which data were averaged over time intervals larger than 5 y, those in which sizes differing by more than one order of magnitude were averaged, and those lacking reliable measurements or comparable restored and reference conditions were discarded, leaving 124 articles (see Text S2). Reference wetlands were usually adjacent to restored or created wetlands, although in some cases they were separated by several kilometers (maximum distance found was ~100 km). In all cases, restored or created wetlands were of the same wetland hydrogeomorphic type [17] as reference wetlands with which they were compared. From the selected articles, six were carried out on experimental wetlands, the rest were carried out on wetland restoration or creation projects. Studies either described measurements at a known age after wetlands were restored or created, or a chronosequence of the progression during the wetland restoration process. Restored and created wetlands were located in 12 countries and totaled >21,294 ha in area and reference wetlands >19,694 ha. The exact total area is not known because it was not reported in 23 out of the 124 selected studies.

Data Extraction

Measurements of structural components and biogeochemical processes were extracted from the main text, tables, and figures of the articles. When abundance of one species was measured at different life stages, only the adult abundance of each species was selected. Variables describing hydrological structure, biological structure, element storage and cycling, and organic matter accumulation were classified as structural components or biogeochemical processes according to wetland functions described by Smith et al. [42], and as ecosystem services described in the Millennium Ecosystem Assessment (MEA) (organic matter accumulation was sometimes designated as “soil formation” in the MEA but not in other soil science references) [49].

Element storage and cycling variables measured processes (mineralization or denitrification) and concentration of elements in different pools (total content in soil, organic content in soil, or content in roots), which suggest how nutrients are moving between pools through biotic and abiotic processes (Tables S1 and S2). The studies presented enough data points to plot recovery of storage of carbon, nitrogen, and phosphorus.

Response Ratio Calculation

To standardize and compare data, we used standard response ratios used in meta-analysis, $\ln(X_{rest}+1/X_{ref}+1)$ [3], where X_{rest} is the value of the measured variable in the restored or created wetland and X_{ref} is the value of the measured variable in the reference wetland. To avoid the value “0” in the natural logarithm of the equation, “1” was added to both values in restored or created and reference wetlands. The effect of adding “1” to the values in the response ratio equation has been demonstrated to have little effect on conclusions [50]. The effect size was not weighted because variance was reported for only 64% of the variables. Differences between weighted and unweighted meta-analysis statistics are generally small [7].

As variables depicting structural components and biogeochemical processes in restored or created wetlands converged to values in the reference wetlands, recovery of function was generally enhanced. But for some variables, such as soil bulk density [51,52], or proportions of exotic species [53,54], higher values are associated with lower levels of wetland recovery. In some cases, the specific context of a study made variables negative for recovery of a particular restored wetland, e.g., the presence of woody species where none had occurred in the reference wetlands [55,56]. In these cases (11% of the collected variables), we changed the sign to reverse the value of the response ratio.

Data Classification

For each variable we recorded the age of the restored or created wetland, the wetland hydrogeomorphic type, the number of restored or created and reference wetlands considered in a given study, the size (ha) of the restored or created and reference wetlands, the initial condition (restored or created), the geographic location, and the climate. Most data (49%) were from wetlands that had been restored or created for less than 5 y (Figure S1). If data from several wetlands of different sizes were averaged in the study, then we also averaged the sizes for our analysis. The geographic location was registered as the latitude and longitude in degrees of the center of the wetland or group of wetlands. The climate was classified according to the last revision of the Köppen-Geiger climate classification [57]. We used the name humid temperate climate for Cf climate, humid cold climate for Df climate, seasonal temperate climate for Cs climate (with dry summer), and seasonal tropical for A climates. Two of our sampled studies were done in seasonal temperate climate with dry winter (Köppen-Geiger climate classification Cw), and were not considered in our climate study. Wetland hydrogeomorphic type was classified according to Brinson [58] and Smith et al. [42] as depressionnal, riverine, tidal, peatland, lacustrine, and seeping slope. Only three studies were on lacustrine wetlands and one on seeping slope wetlands, so these types were not considered in our study of differences among wetland types.

In studies where more than one wetland was studied and data were available for each individual wetland, data were collected for each wetland. In 27 studies, more than one wetland was compared with the same reference wetland, and in 11 studies, restored or created wetlands were compared with more than one reference wetland. All studies where more reference rather than restored or created wetlands were studied provided only averaged data for both groups of wetlands. We calculated contingency tables between the wetland size, the initial conditions (created versus restored), and the covariates included in the environmental setting section (climate and wetland hydrogeomorphic type), using contingency coefficients (C), to test for independence between them. Wetland type showed relevant degrees of association with the climate ($C=0.63$) and wetland size ($C=0.58$), the rest of variables had coefficients below 0.5, indicating low degree of association. These associations may be explained by the influence of the climate on wetland types, e.g., peatlands are usually associated to cold climates, and mangroves to tropical climates. Also, peatlands usually extend over vast surfaces (hundreds or thousands of hectares) and depressionnal wetlands are usually small basins (less than 10 ha or few tens of hectares).

Statistical Analysis

Because data were non-normally distributed (according to the Kolmogorov-Smirnoff test for normality), we used Wilcoxon signed rank tests to test for significant deviations from zero (no

difference from reference conditions) for each estimated mean of the response ratios for variables at each age interval of a restored or created wetland. To test for differences between the same variable measured under two different environmental settings at a given recovery time, we used Kruskal-Wallis tests.

Chronosequences Plotting

To plot the temporal trends, the mean values and the standard error of each variable with every age class of 5 y (0–4.9, 5–9.9, etc) were used. The criterion for a mean for a certain age class to be used in the plot was that it must have been derived from at least nine different data points obtained from at least two different studies. When this criterion was not fulfilled, the mean values and standard error of age classes of 10 y (e.g., 10–19.9, or 20–30) were used. Temporal trend lines were fitted when enough data to calculate means for two or more age classes were available.

Supporting Information

Figure S1 Distribution of wetland sizes across wetland ages for the 654 restored and created wetlands considered in the study.

(TIF)

Figure S2 Chronosequences for the storage and cycling of carbon and nitrogen (A), and for the accumulation of organic matter in soils (B).

(TIF)

Figure S3 Chronosequences for biogeochemical processes (A) and for biological structures (B) under contrasting initial conditions (restored wetlands versus wetlands created de novo in dry lands).

(TIF)

Figure S4 Evolution of the response ratios of restored or created wetlands at successive size categories for wetlands between 5 y to 15 y after restoration or creation.

(TIF)

Table S1 Variables measuring structural components.

(DOC)

Table S2 Variables measuring biogeochemical processes.

(DOC)

Table S3 Statistical significance of differences between the means of the response ratios in restored or created versus reference wetlands.

(DOC)

Table S4 Statistical significance of differences between the means of the response ratios in restored or created versus reference wetlands at each size interval.

(DOC)

Table S5 Statistical significance of differences between the response ratios in restored or created wetlands under different environmental settings.

(DOC)

Text S1 Wetland restoration investment and carbon storage calculation.

(DOC)

Text S2 References used in the meta-analysis.

(DOC)

Acknowledgments

We acknowledge the comments from Joy B. Zedler, Oswald Schmidt, J.M. Rey Benayas, William E. Dietrich, Andrea Swei, and one anonymous reviewer on earlier versions of the manuscript, and the collaboration of Maribel Vara Rodriguez.

References

- Costanza R, d'Arge R, deGroot R, Farber S, Grasso M, et al. (1997) The value of the world's ecosystem services and natural capital. *Nature* 387: 253–260.
- Mitsch WJ, Gosselink JG (2007) *Wetlands*. Hoboken (New Jersey): John Wiley & Sons, Inc. pp 177–183.
- Millennium Ecosystem Assessment (2005) *Ecosystems and human well-being: wetlands and water*. Washington (D.C.): World Resources Institute.
- Palmer MA (2009) Reforming watershed restoration: science in need of application and applications in need of science. *Estuaries Coasts* 32: 1–17.
- Matthews JW, Endress AG (2008) Performance criteria, compliance success, and vegetation development in compensatory mitigation wetlands. *Environ Manage* 41: 130–141.
- Zedler JB, West JM (2008) Declining diversity in natural and restored salt marshes: a 30-year study of Tijuana Estuary. *Restor Ecol* 16: 249–262.
- Rey Benayas JM, Newton AC, Diaz A, Bullock JM (2009) Enhancement of biodiversity and ecosystem services by ecological restoration: a meta-analysis. *Science* 325: 1121–1124.
- Borja A, Dauer DM, Elliott M, Simenstad CA (2010) Medium- and long-term recovery of estuarine and coastal ecosystems: patterns, rates and restoration effectiveness. *Estuaries Coasts* 33: 1249–1260.
- Copeland C (2010) *Wetlands: an overview of issues*. Washington (D.C.): Congressional Research Service. RL33483.
- Palmer MA, Filoso S (2009) Restoration of ecosystem services for environmental markets. *Science* 325: 575–576.
- Zedler JB (2000) Progress in wetland restoration ecology. *Trends Ecol Evol* 15: 402–407.
- Jones HP, Schmitz OJ (2009) Rapid recovery of damaged ecosystems. *Plos One* 4: e5653. doi:10.1371/journal.pone.0005653.
- Rustad LE, Campbell JL, Marion GM, Norby RJ, Mitchell MJ, et al. (2001) A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming. *Oecologia* 126: 543–562.
- Hobbs RJ, Higgs E, Harris JA (2009) Novel ecosystems: implications for conservation and restoration. *Trends Ecol Evol* 24: 599–605.
- Suding KN, Gross KL, Houseman GR (2004) Alternative states and positive feedbacks in restoration ecology. *Trends Ecol Evol* 19: 46–53.
- Suding KN, Hobbs RJ (2009) Threshold models in restoration and conservation: a developing framework. *Trends Ecol Evol* 24: 271–279.
- Ramsar Convention Secretariat (2006) *The Ramsar Convention Manual: a guide to the Convention on Wetlands (Ramsar, Iran, 1971)*, 4th edition. Gland, Switzerland: Ramsar Convention Secretariat.
- Cole CA (1999) Ecological theory and its role in the rehabilitation of wetlands. Streever WR, ed. *An international perspective on wetland rehabilitation*. Dordrecht, The Netherlands: Kluwer Academic Publishers. pp 232–243.
- Hunt RJ, Walker J, Krabbenhoft D (1999) Characterizing hydrology and the importance of ground-water discharge in natural and constructed wetlands. *Wetlands* 19: 458–472.
- Ahn C, Dee S (2011) Early development of plant community in a created mitigation wetland as affected by introduced hydrologic design elements. *Ecol Eng* 37: 1324–1333.
- Kumar JLG, Zhao YQ (2011) A review on numerous modeling approaches for effective, economical and ecological treatment wetlands. *J Environ Manage* 92: 400–406.
- Collins BD, Montgomery DR (2002) Forest development, wood jams, and restoration of floodplain rivers in the Puget Lowland, Washington. *Restor Ecol* 10: 237–247.
- Crooks S, Schutten J, Sheern GD, Pye K, Davy AJ (2002) Drainage and Elevation as Factors in the Restoration of Salt Marsh in Britain. *Restor Ecol* 10: 591–602.
- Lesbarreres D, Fowler MS, Pagano A, Lode T (2010) Recovery of anuran community diversity following habitat replacement. *J Appl Ecol* 47: 148–156.
- Warren RS, Fell PE, Rozsa R, Brawley AH, Orsted AC, et al. (2002) Salt marsh restoration in Connecticut: 20 years of science and management. *Restor Ecol* 10: 497–513.
- Levin LA, Talley TS (2002) Natural and manipulated sources of heterogeneity controlling early faunal development of a salt marsh. *Ecol Appl* 12: 1785–1802.
- Figuerola J, Green AJ, Michot TC (2005) Invertebrate eggs can fly: evidence of waterfowl-mediated gene flow in aquatic invertebrates. *Am Nat* 165: 274–280.
- Badosa A, Frisch D, Arechederra A, Serrano L, Green AJ (2010) Recovery of zooplankton diversity in a restored Mediterranean temporary marsh in Doñana National Park (SW Spain). *Hydrobiologia* 654: 67–82.

Author Contributions

The author(s) have made the following declarations about their contributions: Conceived and designed the experiments: DMM MEP. Analyzed the data: DMM. Wrote the paper: DMM MEP. Performed the meta-analysis: DMM RY. Interpreted and discussed results from the meta-analysis: DMM MEP FAC.

- Meyer CK, Whiles MR (2008) Macroinvertebrate communities in restored and natural Platte River slough wetlands. *J N Am Benthol Soc* 27: 626–639.
- Donath TW, Holzel N, Otte A (2003) The impact of site conditions and seed dispersal on restoration success in alluvial meadows. *Appl Veg Sci* 6: 13–22.
- Matthews JW, Endress AG (2010) Rate of succession in restored wetlands and the role of site context. *Appl Veg Sci* 13: 346–355.
- Zedler JB, Callaway JC (1999) Tracking wetland restoration: do mitigation sites follow desired trajectories? *Restor Ecol* 7: 69–73.
- Knops JMH, Tilman D (2000) Dynamics of soil nitrogen and carbon accumulation for 61 years after agricultural abandonment. *Ecology* 81: 88–98.
- Ballantine K, Schneider R (2009) Fifty-five years of soil development in restored freshwater depressional wetlands. *Ecol Appl* 19: 1467–1480.
- Bechtold JS, Naiman RJ (2009) A quantitative model of soil organic matter accumulation during floodplain primary succession. *Ecosystems* 12: 1352–1368.
- van Groenigen KJ, Six J, Hungate BA, de Graaff MA, van Breemen N, et al. (2006) Element interactions limit soil carbon storage. *Proc Natl Acad Sci U S A* 103: 6571–6574.
- Lawrence D, Schlesinger WH (2001) Changes in soil phosphorus during 200 years of shifting cultivation in Indonesia. *Ecology* 82: 2769–2780.
- Smil V (2000) Phosphorus in the environment: natural flows and human interferences. *Annu Rev Energ Env* 25: 53–88.
- Korfel CA, Mitsch WJ, Hetherington TE, Mack JJ (2010) Hydrology, physiochemistry, and amphibians in natural and created vernal pool wetlands. *Restor Ecol* 18: 843–854.
- Maurer BA (2006) Ecological restoration from a macroscopic perspective. Falk DA, Palmer MA, Zedler JD, eds. *Foundations of restoration ecology*. Washington (D.C.): Island Press.
- Whittaker RJ (2006) *Island biogeography: ecology, evolution, and conservation*. Oxford, UK: Oxford University Press.
- Anderson-Teixeira KJ, Vitousek PM, Brown JH (2008) Amplified temperature dependence in ecosystems developing on the lava flows of Mauna Loa, Hawai'i. *Proc Natl Acad Sci U S A* 105: 228–233.
- Brinson MM (1993) A hydrogeomorphic classification for wetlands. Washington (D.C.): US Army Corps of Engineers. WRP-DE-4. 103 p.
- Craft C, Megonigal P, Broome S, Stevenson J, Freese R, et al. (2003) The pace of ecosystem development of constructed *Spartina alterniflora* marshes. *Ecol Appl* 13: 1417–1432.
- Dolinar N, Gaberscik A (2010) Mycorrhizal colonization and growth of *Phragmites australis* in an intermittent wetland. *Aquat Bot* 93: 93–98.
- Kardol P, Wardle DA (2010) How understanding aboveground belowground linkages can assist restoration ecology. *Trends Ecol Evol* 25: 670–679.
- De Deyn GB, Cornelissen JHC, Bardgett RD (2008) Plant functional traits and soil carbon sequestration in contrasting biomes. *Ecol Lett* 11: 516–531.
- Race MS, Fonseca MS (1996) Fixing compensatory mitigation: what will it take? *Ecol Appl* 6: 94–101.
- Smith RD, Ammann A, Bartoldus C, Brinson MM (1995) An approach for assessing wetland functions using hydrogeomorphic classification, reference wetlands, and functional indices. Washington (D.C.): US Army Corps of Engineers.
- Scheiner SM, Gurevitch J (2001) *Design and analysis of ecological experiments*. Cary/North Carolina: Oxford University Press, Incorporated.
- Cardinale BJ, Srivastava DS, Duffy JE, Wright JP, Downing AL, et al. (2006) Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature* 443: 989–992.
- Marvier M, McCreedy C, Regetz J, Kareiva P (2007) A meta-analysis of effects of Bt cotton and maize on nontarget invertebrates. *Science* 316: 1475–1477.
- Craft C, Broome S, Campbell C (2002) Fifteen years of vegetation and soil development after brackish-water marsh creation. *Restor Ecol* 10: 248–258.
- Ballantine K, Schneider R (2009) Fifty-five years of soil development in restored freshwater depressional wetlands. *Ecol Appl* 19: 1467–1480.
- Gutrich JJ, Taylor KJ, Fennessy MS (2009) Restoration of vegetation communities of created depressional marshes in Ohio and Colorado (USA): the importance of initial effort for mitigation success. *Ecol Eng* 35: 351–368.
- Parikh A, Gale N (1998) Vegetation monitoring of created dune swale wetlands, Vandenberg Air Force Base, California. *Restor Ecol* 6: 83–93.
- Zampella RA, Laidig KJ (2003) Functional equivalency of natural and excavated coastal plain ponds. *Wetlands* 23: 860–876.
- Kottek M, Griesser J, Beck C, Rudolf B, Rubel F (2006) World Map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift* 15: 259–263.